

CORE PEDIATRIC AND ADOLESCENT MEDICINE

9 Month Handouts

212 Calef Highway, Epping NH 03042

603-693-2100

Core Pediatrics Vaccination and Screening Schedule

WCC visit	Recommended Vaccines	Recommended Screening Tests
Birth/Hospital	Hepatitis B	Hearing screening, State Metabolic screening, Congenital Heart Disease screening
2 Week	Hepatitis B (if not given in hospital)	
6 Week	Pentacel*, Hepatitis B, Prevnar, Rotateq	
4 Month	Pentacel*, Prevnar, Rotateq (Hepatitis B if not received at birth)	
6 Month	Pentacel*, Hepatitis B, Prevnar, Rotateq	
9 Month		Developmental screening (ASQ) and Hemoglobin (optional)
12 Month	MMR*, Varivax, Hepatitis A	Photoscreening, Lead, Hemoglobin, Fluoride
15 Month	Pentacel*, Prevnar	
18 Month	Hepatitis A	Developmental screening (ASQ and MCHAT)
2 Year	Hepatitis A (if not given at 18 months)	Developmental screening (MCHAT), Lead, Hemoglobin, Photoscreening, Fluoride
2 ½ Year		Developmental screening (ASQ)
3 Year		Photoscreening
4 Year	Kinrix*, Proquad*	Vision and Hearing
5 -6 Year		Vision and Hearing
7 Year		
8 Year		Vision and Hearing
9 Year		
10 Year	TDaP	Vision and Hearing, Cholesterol
11 Year	Menactra, HPV#	
12 Year	(HPV see schedule below)	Depression screening (PHQ-9 or Ipad), Vision
13 Year		Depression screening (PHQ-9 or Ipad)
14 Year		Depression screening (PHQ-9 or Ipad), Urine STI screening
15 Year		Depression screening (PHQ-9 or Ipad), Urine STI screening, Vision
16 Year	Menactra, Men B-Bexsero	Depression screening (PHQ-9 or Ipad), Urine STI screening
17 Year		Depression screening (PHQ-9 or Ipad), Urine STI screening
18 Year		Depression Screening (PHQ-9 or Ipad), Urine STI screening, Cholesterol, HIV, Vision (optional)
19-20 Year	Tdap Booster	Depression screening (PHQ-9 or Ipad), Urine STI screening
21 Year		Depression screening (PHQ-9 or Ipad), Urine STI screening, Pap smear

If clinically indicated:

Influenza (6 months plus), Pneumovax 23, Prevnar 13, PPD, Typhoid, Men B


#HPV Dosing Interval:

- If received first dose **before 15th birthday** : 0 and 6 months
- If received first dose **after 15th birthday**: 0, 2 and 6 months

***Combination Vaccines**

Pentacel- DTaP, Hib, IPV
MMR- Measles, Mumps, Rubella
Kinrix- DTaP, IPV
Proquad- MMR, Varicella
Pediarix- DTaP, Hepatitis B, IPV

6 to 12 Months



6 TO 12 MONTHS

Safety for Your Child

Did you know that hundreds of children younger than 1 year die every year in the United States because of injuries — most of which can be prevented?

Often, injuries happen because parents are not aware of what their children can do. Your child is a fast learner and will suddenly be able to *roll over*, *crawl*, *sit*, and *stand*. Your child may *climb* before walking, or *walk* with support months before you expect. Your child will *grasp* at almost anything and reach things he or she could not reach before.

Falls

Because of your child's new abilities, he or she will fall often. Protect your child from injury. **Use gates on stairways and doors. Install operable window guards** on all windows above the first floor. **Remove sharp-edged or hard furniture** from the room where your child plays.

Do not use a baby walker. Your child may tip it over, fall out of it, or fall down the stairs in it. Baby walkers allow children to get to places where they can pull hot foods or heavy objects down on themselves.


If your child has a serious fall or does not act normally after a fall, call your doctor.

Burns

At 6 to 12 months children grab at everything. **NEVER** leave cups of hot coffee on tables or counter edges. **And NEVER carry hot liquids or food near your child or while holding your child.** He or she could get burned. Also, if your child is left to crawl or walk around stoves, wall or floor heaters, or other hot appliances, he or she is likely to get burned. **A safer place for your child** while you are cooking, eating, or unable to provide your full attention is the **playpen, crib, or stationary activity center, or buckled into a high chair.**

If your child does get burned, put cold water on the burned area immediately. Keep the burned area in cold water for a few minutes to cool it off. Then cover the burn loosely with a dry bandage or clean cloth. Call your doctor for all burns. To protect your child from tap water scalds, the hottest temperature at the faucet should be no more than 120°F. In many cases you can adjust your water heater.

Make sure you have a working smoke alarm on every level of your home, especially in furnace and sleeping areas. Test the alarms every month. It is best to use smoke alarms that use long-life batteries, but if you do not, change the batteries at least once a year.



(over)

American Academy of Pediatrics

DEDICATED TO THE HEALTH OF ALL CHILDREN™



Drowning

At this age your child loves to play in water. Empty all the water from a bathtub, pail, or any container of water immediately after use. Keep the door to the bathroom closed. **NEVER leave your child alone in or near a bathtub, pail of water, wading or swimming pool, or any other water, even for a moment.** Drowning can happen in less than 2 inches of water. Knowing how to swim does NOT mean your child is safe in or near water. Stay within an arm's length of your child around water.

If you have a swimming pool, now is the time to **install a fence** that separates the house from the pool. The pool should be fenced in on all 4 sides. Most children drown when they wander out of the house and fall into a pool that is not fenced off from the house. Be prepared — install a fence around your pool now, before your child begins to walk!



Poisoning and Choking

Your child will explore the world by *putting anything and everything into his or her mouth*. NEVER leave small objects or balloons in your child's reach, even for a moment. Don't feed your child hard pieces of food such as hot dogs, raw carrots, grapes, peanuts, or popcorn. Cut all of his or her food into thin slices to prevent choking.

Be prepared if your child starts to choke. Learn how to save the life of a choking child. Ask your doctor to recommend the steps you need to take.

Children will put everything into their mouths, even if it doesn't taste good. Many ordinary things in your house **can be poisonous** to your child. Be sure to keep household products such as cleaners, chemicals, and medicines up, up, and away, completely out of sight and reach. Never store lye drain cleaners in your home. **Use safety latches or locks** on drawers and cupboards. Remember, your child doesn't understand or remember "no" while exploring.



If your child does eat something that could be poisonous, call the Poison Help number at 1-800-222-1222 immediately. Do not make your child vomit.

Strangulation and Suffocation

Place your baby's crib away from windows. **Cords from window blinds and draperies can strangle your child.** Use cordless window coverings, or if this is not possible, tie cords high and out of reach. Do not knot cords together.

Plastic wrappers and bags form a tight seal if placed over the mouth and nose and may suffocate your child. Keep them away from your child.

And Remember Car Safety

Car crashes are a **great danger** to your child's life and health. Most injuries and deaths caused by car crashes **can be prevented** by the use of car safety seats **EVERY TIME** your child is in the car. All infants and toddlers should ride in a rear-facing car safety seat until they are at least 2 years of age or until they reach the highest weight or height allowed by their car safety seat's manufacturer. A rear-facing car safety seat should **NEVER** be placed in front of a passenger air bag. Your child, besides being much safer in a car safety seat, will behave better so you can pay attention



to your driving. **The safest place for all infants and children to ride is in the back seat.**

Do not leave your child alone in a car. Keep vehicles and their trunks locked. Children who are left in a car can die of heat stroke because temperatures can reach deadly levels in minutes.

Remember, the biggest threat to your child's life and health is an injury.

From Your Doctor

The information in this publication should not be used as a substitute for the medical care and advice of your pediatrician. There may be variations in treatment that your pediatrician may recommend based on the individual facts and circumstances.

Activities for Infants 8-12 Months Old



<p>Let your baby feed himself. This gives your baby practice picking up small objects (cereal, cooked peas) and also gives him experience with textures in his hands and mouth. Soon your baby will be able to finger feed an entire meal.</p>	<p>Your baby will be interested in banging objects to make noise. Give your baby blocks to bang, rattles to shake, or wooden spoons to bang on containers. Show your baby how to bang objects together.</p>	<p>A good pastime is putting objects in and out of containers. Give your baby plastic containers with large beads or blocks. Your baby may enjoy putting socks in and out of the sock drawer or small cartons (Jell-O, tuna or soup cans) on and off shelves.</p>	<p>Mirrors are exciting at this age. Let your baby pat and poke at herself in the mirror. Smile and make faces together in the mirror.</p>	<p>Your baby will begin using his index fingers to poke. Let your baby poke at a play telephone or busy box. Your baby will want to poke at faces. Name the body parts as your baby touches your face.</p>
<p>Put toys on a sofa or sturdy table so that your baby can practice standing while playing with the toys.</p>	<p>Find a big box that your baby can crawl in and out of. Stay close by and talk to your baby about what she is doing. "You went in! Now you are out!"</p>	<p>Read baby books or colorful magazines by pointing and telling your baby what is in the picture. Let your baby pat pictures in the book.</p>	<p>Play hide-and-seek games with objects. Let your baby see you hide an object under a blanket, diaper, or pillow. If your baby doesn't uncover the object, just cover part of it. Help your baby find the object.</p>	<p>Play ball games. Roll a ball to your baby. Help your baby, or have a partner help him roll the ball back to you. Your baby may even throw the ball, so beach balls or Nerf balls are great for this game.</p>
<p>Turn on a radio or stereo. Hold your baby in a standing position and let your baby bounce and dance. If your baby can stand with a little support, hold her hands and dance like partners.</p>	<p>Play imitation games like Peek-a-boo and So Big. Show pleasure at your baby's imitations of movements and sounds. Babies enjoy playing the same games over and over.</p>	<p>Let your baby play with plastic measuring cups, cups with handles, sieves and strainers, sponges, and balls that float in the bathtub. Bath time is a great learning time.</p>	<p>Play Pat-a-Cake with your baby. Clap his hands together or take turns. Wait and see if your baby signals you to start the game again. Try the game using blocks or spoons to clap and bang with.</p>	<p>Your baby will play more with different sounds like "la-la" and "da-da." Copy the sounds your baby makes. Add a new one and see if your baby tries it, too. Enjoy your baby's early attempts at talking.</p>
<p>Make a simple puzzle for your baby by putting blocks or Ping-Pong balls inside a muffin pan or egg carton.</p>	<p>You can make a simple toy by cutting a round hole in the plastic lid of a coffee can. Give your baby wooden clothes pins or Ping-Pong balls to drop inside.</p>	<p>Say "hi" and wave when entering a room with your baby. Encourage your baby to imitate. Help your baby wave to greet others. Waving "hi" and "bye" are early gestures.</p>	<p>Let your baby make choices. Offer two toys or foods and see which one your baby picks. Encourage your baby to reach or point to the chosen object. Babies have definite likes and dislikes!</p>	<p>New places and people are good experiences for your baby, but these can be frightening. Let your baby watch and listen and move at her own speed. Go slowly. Your baby will tell you when she is ready for more.</p>

Activities for Infants 8 -12 Months Old



<p>Let your baby feed himself. This gives your baby practice picking up small objects (cereal, cooked peas) and also gives him experience with textures in his hands and mouth. Soon your baby will be able to finger feed an entire meal.</p>	<p>Your baby will be interested in banging objects to make noise. Give your baby blocks to bang, rattles to shake, or wooden spoons to bang on containers. Show your baby how to bang objects together.</p>	<p>A good pastime is putting objects in and out of containers. Give your baby plastic containers with large beads or blocks. Your baby may enjoy putting socks in and out of the sock drawer or small cartons (Jell-O, tuna or soup cans) on and off shelves.</p>	<p>Mirrors are exciting at this age. Let your baby pat and poke at herself in the mirror. Smile and make faces together in the mirror.</p>	<p>Your baby will begin using his index fingers to poke. Let your baby poke at a play telephone or busy box. Your baby will want to poke at faces. Name the body parts as your baby touches your face.</p>
<p>Put toys on a sofa or sturdy table so that your baby can practice standing while playing with the toys.</p>	<p>Find a big box that your baby can crawl in and out of. Stay close by and talk to your baby about what she is doing. "You went in! Now you are out!"</p>	<p>Read baby books or colorful magazines by pointing and telling your baby what is in the picture. Let your baby pat pictures in the book.</p>	<p>Play hide-and-seek games with objects. Let your baby see you hide an object under a blanket, diaper, or pillow. If your baby doesn't uncover the object, just cover part of it. Help your baby find the object.</p>	<p>Play ball games. Roll a ball to your baby. Help your baby, or have a partner help him roll the ball back to you. Your baby may even throw the ball, so beach balls or Nerf balls are great for this game.</p>
<p>Turn on a radio or stereo. Hold your baby in a standing position and let your baby bounce and dance. If your baby can stand with a little support, hold her hands and dance like partners.</p>	<p>Play imitation games like Peek-a-boo and So Big. Show pleasure at your baby's imitations of movements and sounds. Babies enjoy playing the same games over and over.</p>	<p>Let your baby play with plastic measuring cups, cups with handles, sieves and strainers, sponges, and balls that float in the bathtub. Bath time is a great learning time.</p>	<p>Play Pat-a-Cake with your baby. Clap his hands together or take turns. Wait and see if your baby signals you to start the game again. Try the game using blocks or spoons to clap and bang with.</p>	<p>Your baby will play more with different sounds like "la-la" and "da-da." Copy the sounds your baby makes. Add a new one and see if your baby tries it, too. Enjoy your baby's early attempts at talking.</p>
<p>Make a simple puzzle for your baby by putting blocks or Ping-Pong balls inside a muffin pan or egg carton.</p>	<p>You can make a simple toy by cutting a round hole in the plastic lid of a coffee can. Give your baby wooden clothes pins or Ping-Pong balls to drop inside.</p>	<p>Say "hi" and wave when entering a room with your baby. Encourage your baby to imitate. Help your baby wave to greet others. Waving "hi" and "bye" are early gestures.</p>	<p>Let your baby make choices. Offer two toys or foods and see which one your baby picks. Encourage your baby to reach or point to the chosen object. Babies have definite likes and dislikes!</p>	<p>New places and people are good experiences for your baby, but these can be frightening. Let your baby watch and listen and move at her own speed. Go slowly. Your baby will tell you when she is ready for more.</p>



Bright Futures Parent Handout 9 Month Visit

Here are some suggestions from Bright Futures experts that may be of value to your family.

Your Baby and Family

FAMILY ADAPTATIONS

- Tell your baby in a nice way what to do ("Time to eat"), rather than what not to do.
- Be consistent.
- At this age, sometimes you can change what your baby is doing by offering something else like a favorite toy.
- Do things the way you want your baby to do them—you are your baby's role model.
- Make your home and yard safe so that you do not have to say "No!" often.
- Use "No!" only when your baby is going to get hurt or hurt others.
- Take time for yourself and with your partner.
- Keep in touch with friends and family.
- Invite friends over or join a parent group.
- If you feel alone, we can help with resources.
- Use only mature, trustworthy babysitters.
- If you feel unsafe in your home or have been hurt by someone, let us know; we can help.

Feeding Your Baby

FEEDING ROUTINE

- Be patient with your baby as he learns to eat without help.
- Being messy is normal.
- Give 3 meals and 2–3 snacks each day.
- Vary the thickness and lumpiness of your baby's food.
- Start giving more table foods.
- Give only healthful foods.
- Do not give your baby soft drinks, tea, coffee, and flavored drinks.
- Avoid forcing the baby to eat.
- Babies may say no to a food 10–12 times before they will try it.
- Help your baby to use a cup.

FEEDING ROUTINE

- Continue to breastfeed or bottle-feed until 1 year; do not change to cow's milk.
- Avoid feeding foods that are likely to cause allergy—peanut butter, tree nuts, soy and wheat foods, cow's milk, eggs, fish, and shellfish.

Your Changing and Developing Baby

INFANT INDEPENDENCE

- Keep daily routines for your baby.
- Make the hour before bedtime loving and calm.
- Check on, but do not pick up, the baby if she wakes at night.
- Watch over your baby as she explores inside and outside the home.
- Crying when you leave is normal; stay calm.
- Give the baby balls, toys that roll, blocks, and containers to play with.
- Avoid the use of TV, videos, and computers.
- Show and tell your baby in simple words what you want her to do.
- Avoid scaring or yelling at your baby.
- Help your baby when she needs it.
- Talk, sing, and read daily.

Safety

SAFETY

- Use a rear-facing car safety seat in the back seat in all vehicles.
- Have your child's car safety seat rear-facing until your baby is 2 years of age *or* until she reaches the highest weight or height allowed by the car safety seat's manufacturer.
- Never put your baby in the front seat of a vehicle with a passenger air bag.
- Always wear your own seat belt and do not drive after using alcohol or drugs.
- Empty buckets, pools, and tubs right after you use them.

SAFETY

- Place gates on stairs; do not use a baby walker.
- Do not leave heavy or hot things on tablecloths that your baby could pull over.
- Put barriers around space heaters, and keep electrical cords out of your baby's reach.
- Never leave your baby alone in or near water, even in a bath seat or ring. Be within arm's reach at all times.
- Keep poisons, medications, and cleaning supplies locked up and out of your baby's sight and reach.
- Call Poison Help (1-800-222-1222) if you are worried your child has eaten something harmful.
- Install openable window guards on second-story and higher windows and keep furniture away from windows.
- Never have a gun in the home. If you must have a gun, store it unloaded and locked with the ammunition locked separately from the gun.
- Keep your baby in a high chair or playpen when in the kitchen.

What to Expect at Your Child's 12 Month Visit

We will talk about

- Setting rules and limits for your child
- Creating a calming bedtime routine
- Feeding your child
- Supervising your child
- Caring for your child's teeth

Poison Help: 1-800-222-1222

Child safety seat inspection:
1-866-SEATCHECK; seatcheck.org



American Academy
of Pediatrics



DEDICATED TO THE HEALTH OF ALL CHILDREN™

The recommendations in this publication do not indicate an exclusive course of treatment or serve as a standard of medical care. Variations, taking into account individual circumstances, may be appropriate. Original document included as part of Bright Futures Tool and Resource Kit. Copyright © 2010 American Academy of Pediatrics, Updated 8/11. All Rights Reserved. The American Academy of Pediatrics does not review or endorse any modifications made to this document and in no event shall the AAP be liable for any such changes.